

Case report

Open Access

Compression of the median nerve in the proximal forearm by a giant lipoma: A case report

Sebastian E Valbuena*¹, Greg A O'Toole² and Eric Roulot²

Address: ¹Department of Orthopaedic Surgery & Traumatology, Hospital Interzonal El Cruce, Alta complejidad en red. Florencio Varela, Buenos Aires, Argentina and ²Institut de la Main, Clinique Jouvenet, Paris, France

Email: Sebastian E Valbuena* - valbuena.sebastian@gmail.com; Greg A O'Toole - gregotoole@gmail.com; Eric Roulot - eroulot@free.fr

* Corresponding author

Published: 10 June 2008

Received: 21 February 2008

Journal of Brachial Plexus and Peripheral Nerve Injury 2008, **3**:17 doi:10.1186/1749-7221-3-17

Accepted: 10 June 2008

This article is available from: <http://www.jbppni.com/content/3/1/17>

© 2008 Valbuena et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Compression of the median nerve by a tumour in the elbow and forearm region is rare. We present a case of neuropathy of the median nerve secondary to compression by giant lipoma in the proximal forearm.

Case presentation: A 46-year-old man presented with a six month history of gradually worsening numbness and paresthesia on the palmar aspect of the left thumb and thenar eminence. Clinical examination reveals a hypoaesthesia in the median nerve area of the left index and thumb compared to the contralateral side. Electromyography showed prolonged sensory latency in the distribution of the median nerve corresponding to compression in the region of the pronator teres (pronator syndrome). Radiological investigations were initially reported as normal. Conservative treatment for one month did not result in any improvement. Surgical exploration was performed and a large intermuscular lipoma enveloped the median nerve was found. A complete excision of the tumour was performed. Postoperative reevaluation the X-ray of the elbow was seen to demonstrate a well-circumscribed mass in the anterior aspect of the proximal forearm. At follow-up, 14 months after surgery, the patient noted complete return of the sensation and resolution of the paresthesia.

Conclusion: In case of atypical findings or non frequent localization of nerve compression, clinically interpreted as an idiopathic compression, it is recommended to make a pre-operative complementary Ultrasound or MRI study.

Background

Compression of the median nerve in the elbow and proximal forearm region is much less frequent than within the carpal tunnel [1]. Proximal compression is most commonly the result of anatomic variations with the supracondylar process and Struthers ligament [2], the lacertus fibrosus (bicipital aponeurosis), the pronator teres muscle and the arch of the flexor superficialis most commonly

implicated [3]. With less frequency, anomalous anatomic structures are implicated in compression of the median nerve, these being most commonly, the accessory head of the flexor pollicis longus (Ganzer's muscle) [4], and a persistent median artery [5].

Rarer causes of extrinsic compression of the median nerve such as chronic compartment syndrome [6], partial rup-

ture of the distal biceps insertion [7], and synovial osteochondromatosis at the elbow [8] have also been reported.

The compressive neuropathy of the median nerve secondary to lipoma is not frequent, and has been described principally in the wrist and the hand [9-12]. We present a case of compression of the median nerve in the proximal forearm by a giant lipoma.

Case presentation

A 46-year-old man presented with a six month history of gradually worsening numbness and paresthesia on the palmar aspect of the left thumb and thenar eminence.

Static two point discrimination in the median nerve distribution of the index and thumb showed a hypoaesthesia compared to the contralateral side. Tinel's sign of the wrist and forearm and Phalen's sign were negative. Active forearm pronation against resistance in slight flexion, resisted active forearm supination and resisted active index and middle finger flexion did not elicit pain. Grip strength was equal bilaterally. No masses were detectable on examination of the hand, wrist and forearm. No history of vaccination, viral infection or medication within the previous year was offered.

X-rays of the cervical vertebrae, elbow, forearm and hand were initially reported as normal. However, on postoperative reevaluation the X-ray of the elbow was seen to demonstrate a well-circumscribed mass in the anterior aspect



Figure 1
A lateral elbow X-ray subtly demonstrates a well circumscribed mass in the anterior proximal forearm (arrows).

of the proximal forearm (Figure 1). Electromyography showed prolonged sensory latency in the distribution of the median nerve corresponding to compression in the region of the pronator teres (pronator syndrome).

Conservative treatment (anti-inflammatory medication and a diurnal long-arm splint) for one month did not result in any improvement. Surgical exploration was therefore performed under regional anesthesia and hemostatic tourniquet. The surgical incision began just medial to the biceps tendon and distal to the elbow flexion crease and continued to the mid-forearm between the flexor and extensor muscle masses. The medial antebrachial cutaneous nerve was identified and retracted. The pronator mass and the biceps tendon were identified. An intermuscular mass of adipose tissue was identified just lateral to the superficial head of the pronator teres, the dissection was not difficult but the median nerve was enveloped by the tumour (Figure 2). Microsurgical techniques were used to allow an extracapsular and non-traumatic dissection. A complete excision of the tumour of 8 cm × 6 cm × 3 cm was performed (Figure 4). The median nerve had an hour-glass deformity as a result of its compression (Figure 3). Histopathological examination of the tissue removed at surgery confirmed the presence of well-differentiated mature fat cells (lipoma). There were no neural or neoplastic features.

At follow-up, 14 months after surgery, the patient noted complete return of the sensation and resolution of the paresthesia.




Figure 2
Intraoperative photo showing the median nerve within the intermuscular lipoma.




Figure 3
Surgical area after lipoma excision: the arrow shows the hourglass deformation of the nerve at the level of the compression.

Discussion

Lipomas are benign tumours originating from adipose cells occurring in subcutaneous tissues, intermuscular, intramuscular or paraosteal localizations [9]. Lipomas of more than 5 cm diameter (Giant lipoma) are infrequent



Figure 4
The giant lipoma was excised completely and measured 8 cm × 6 cm × 3 cm.

in the upper limb [10]. In this anatomic location, masses are generally symptomatic at a smaller size.

Nerve compression by a lipoma is uncommon [9,13] but subfascial lipomas are deep tumours and can be a cause of nerve compression [14].

Review of English literature on extrinsic nerve compression by lipoma in the upper limb reveals several reports of compression of the radial nerve (especially the posterior interosseous nerve due to the anatomic relation with the neck of the radius) [8,15-19], and a few cases of ulnar nerve compression in the forearm [11] and the Guyon's canal [20-22]. The involvement of the median nerve was also reported in the brachial plexus area's [23] and particularly in the wrist or the palmar region's [9-12,24-27]. Only one case documented an extrinsic compression of the median nerve in the proximal forearm by a giant lipoma resulting in an anterior interosseous syndrome [28]. To our knowledge, the case presented is the second reported case of compression of the median nerve in the proximal forearm by a giant lipoma.

Cribb et al. [10] documented a series of 10 giant lipomatous tumours (7 lipomas, one neural fibrolipoma and two well differentiated lipoma-like liposarcomas), five cases were in the hand and five cases in the forearm, with signs of median nerve compression in two cases, one in the hand with the location of compression in the second case being unclear. However, in all cases neurovascular structures required mobilisation in order to excise the tumour.

Cribb et al. [10] stressed the importance of a multidisciplinary approach to investigations of giant soft tissue tumours and suggested that an MRI should be routine. In cases where an MRI does not clearly demonstrate a lipoma or in those patients who could not tolerate the investigation, they go on to suggest that a biopsy be performed.

Johnson et al. [29] demonstrated that soft tissue masses of greater than five cm in diameter should be considered malignant unless proven otherwise.

Marginal resection with conservation of the neurovascular structures is the procedure of choice for lipomas, and a more aggressive surgery is required in case of malign tumour. In our case the diagnosis of tumour compression, despite of the size of the lipoma, was made intra-operatively as we had considered the atypical clinical findings to be the result of idiopathic compression of the median nerve in the elbow region.

Ultrasound is an excellent diagnostic study, especially for deeply sited masses and can be used such cases. MRI however, provides more information of tumour type and of

anatomic relations and is therefore preferable for diagnostic precision and pre-operative planning.

Conclusion

Extrinsic median nerve compression by a tumour is rare. However, in case of atypical findings or non frequent localization of nerve compression, clinically interpreted as an idiopathic compression, it is recommended to make a pre-operative complementary Ultrasound or MRI study.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SEV, GAO, and ER conceived the case report and interpreted the data, SEV performed all pertinent literature review on the subject and drafted the manuscript, ER performed the patient's surgery and collected the clinical data, SEV assisted to ER in the surgery, GAO helped to draft the manuscript.

All authors read and approved the final manuscript.

Acknowledgements

No financial support was provided in any form to the authors of this manuscript.

References

- Gross PT, Jones HR Jr: **Proximal median neuropathies: electromyographic and clinical correlation.** *Muscle Nerve* 1992, **15**:390-395.
- al-Qattan MM, Husband JB: **Median nerve compression by the supracondylar process: a case report.** *J Hand Surg [Br]* 1991, **16**(1):101-103.
- Johnson RK, Spinner M, Shrewsbury MM: **Median nerve entrapment syndrome in the proximal forearm.** *J Hand Surg [Am]* 1979, **4**:48-51.
- Dellon AL, Mackinnon SE: **Musculoaponeurotic variations along the course of the median nerve in the proximal forearm.** *J Hand Surg [Br]* 1987, **12**(3):359-363.
- Jones NF, Ming NL: **Persistent median artery as a cause of pronator syndrome.** *J Hand Surg [Am]* 1988, **13**:728-32.
- Kutz JE, Singer R, Lindsay M: **Chronic exertional compartment syndrome of the forearm: a case report.** *J Hand Surg [Am]* 1985, **10**(2):302-304.
- Foxworthy M, Kinninmonth AWW: **Median nerve compression in the proximal forearm as a complication of partial rupture of the distal biceps brachii tendon.** *J Hand Surg [Br]* 1992, **17**:515-517.
- Nogueira A, Alcelay O, Pena C, Sarasua JG, Madrigal B: **Synovial osteochondromatosis at the elbow producing ulnar and median nerve palsy. Case report and review of the literature.** *Chir Main* 1999, **18**:108-114.
- Leffert RD: **Lipoma of the upper extremity.** *J Bone Joint Surg Am* 1972, **54**:1262-1266.
- Cribb GL, Cool WP, Ford DJ, Mangham DC: **Giant lipomatous tumors of the hand and forearm.** *J Hand Surg [Br]* 2005, **30**(5):509-512.
- Flores LP, Carneiro JZ: **Peripheral nerve compression secondary to adjacent lipomas.** *Surg Neurol* 2007, **67**:258-263.
- Babins DM, Lubahn JD: **Palmar lipomas associated with compression of the median nerve.** *J Bone Joint Surg Am* 1994, **76**(9):1360-1362.
- Phalen GS, Kendrick JI, Rodriguez JM: **Lipomas of the upper extremity. A series of fifteen tumours in the hand and six tumours causing nerve compression.** *Am J Surg* 1971, **121**:298-306.
- Barber KW, Bianco AJ Jr, Soule ED, McCarty CS: **Benign extraneural soft tissues tumours of the extremities causing compression of nerves.** *J Bone Joint Surg Am* 1962, **44**:98-104.
- Bieber EJ, Moore JR, Weiland AJ: **Lipomas compressing the radial nerve at the elbow.** *J Hand Surg [Am]* 1986, **11**(4):533-535.
- Fitzgerald A, Anderson WW, Hooper G: **Posterior interosseous nerve palsy due to parosteal lipoma.** *J Hand Surg [Br]* 2002, **27**:535-537.
- Lidor C, Lotem M, Hallel T: **Parosteal lipoma of the proximal radius: a report of 5 cases.** *J Hand Surg [Am]* 1992, **17**:1095-1097.
- Nishida J, Shimamura T, Ehara S, Shiraishi H, Sato T, Abe M: **Posterior interosseous nerve palsy caused by parosteal lipoma of proximal radius.** *Skeletal Radiol* 1998, **27**:375-379.
- Werner CO: **Radial nerve paralysis and tumor.** *Clin Orthop Relat Res* 1991, **268**:223-225.
- Bui-Mansfield LT, Williamson M, Wheeler DT, Johnstone F: **Guyon's canal lipoma causing ulnar neuropathy.** *AJR Am J Roentgenol* 2002, **178**:1458.
- Galeano M, Colonna M, Risitano G: **Ulnar tunnel syndrome secondary to lipoma of the hypothenar region.** *Ann Plast Surg* 2001, **46**:83-84.
- Zahrawi F: **Acute compression ulnar neuropathy at Guyon's canal resulting from lipoma.** *J Hand Surg [Am]* 1984, **9**(2):238-240.
- Weinzweig N, Browne EZ Jr: **Infraclavicular median nerve compression caused by a lipoma.** *Orthopedics* 1988, **11**:1077-1078.
- Oster LH, Blair WF, Steyers CM: **Large lipomas in the deep palmar space.** *J Hand Surg [Am]* 1989, **14**(4):700-704.
- Schmitz RL, Keeley JL: **Lipomas of the hand.** *Surgery* 1957, **42**:696-700.
- Sonoda H, Takasita M, Taira H, Higashi T, Tsumura H: **Carpal tunnel syndrome and trigger wrist caused by a lipoma arising from flexor tenosynovium: a case report.** *J Hand Surg [Am]* 2002, **27**(6):1056-1058.
- Kremchek TE, Kremchek EJ: **Carpal tunnel syndrome caused by flexor tendon sheath lipoma.** *Orthop Rev* 1988, **17**:1083-1085.
- Higgs PH, Young VL, Schuster R, Weeks PM: **Giant lipomas of the hand and forearm.** *South Med J* 1993, **86**:887-890.
- Johnson CJ, Pynsent PB, Grimer RJ: **Clinical features of soft tissue sarcomas.** *Ann R Coll Surg Engl* 2001, **83**:203-205.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

